

Original Article

Influence of Temperature and Relative Humidity on the Culture of Fungi Isolated from Rotting of Sweet Potato (*Ipomoea batatas* (L.) Lam.)

Bello, I. M^{*1}., Suberu, H. A¹., Kuta, F. A². and Daudu, O. A. Y¹.

¹Department of Biological Sciences, Federal University of Technology, Minna, Nigeria

²Department of Microbiology, Federal University of Technology, Minna, Nigeria

Submitted: March 08, 2012; Accepted: Nov. 12, 2012; Published: Dec. 17, 2012.

ABSTRACT

The influence of temperature and relative humidity on the culture of species of fungi isolated from rotting sweet potato tubers for delineation of physical control measure was investigated in this study. The fungi were cultured at five temperature regimens: 27±3°C (i.e., room temperature, 'Control'), 35, 40, 45 and 50°C; and six Relative Humidity (R.H.) levels: 32, 50, 74, 85, 97 and 100%, for a period of eight days. The growth of the fungi increased till 40°C, after which significant (P<0.05) drop in growth was recorded. However, significant (P>0.05) differences in growth of the fungi were limited to the first three temperature regimens tested. With respect to relative humidity, all the five fungi tested recorded their least growth at the lowest relative humidity (i.e., 32.50%), *Aspergillus niger*, *Rhizopus oryzae* and *Mucor racemosus* achieved their maximum growth at 50% R.H. (i.e., 46±28.07, 53.38±23.37 and 59.63±28.09mm, respectively); while *Fusarium oxysporum* and *Scopulariopsis brevicaulis* grew optimally at 85% R.H. (56.43±28.83 and 28.38±18.02mm, respectively). These results suggest significant shifts in the optimal temperature and R.H. for the growth of some of the fungi. The findings of this study should guide the development of better storage facilities for potato tubers, through the modification of micro-climatic conditions.

Key words: Fungi, Growth, Relative Humidity, and Temperature

***Corresponding Author:** ibellomohammed@yahoo.com

INTRODUCTION

Sweet potato (*Ipomoea batatas*) is one of the most important food crops, yet only 5% of its world production is contributed by Africa (FAO, 1992). A major limitation to optimum production of sweet potato in Africa in general and Nigeria in particular, has been the huge postharvest losses often associated with the tubers under storage conditions (George, 1988; Abubakar, 1990). In Kenya, postharvest soft-rotting of sweet potato

tubers was rated as a major production constraint (Mutuura *et al.*, 1992). According to Kihurani *et al.* (2008), certain storage environmental conditions are known to directly or indirectly predispose sweet potato tubers to postharvest microbial deterioration. Walker (1998) observed that the progression of sweet potato soft-rot diseases was fastest when a high humidity was combined with a temperature of 80°F. For example, the optimum temperature for the growth of *Erwinia* sp, a major causal pathogen of potato soft-rot, has been put at

between 85° and 120°F, with the maximum been slightly over 100°F.

Similar observations have been made with respect to the influence of temperature and R.H. on soft-rot microbial pathogenic activities in other crops. Bhat *et al.* (2010a&b) observed higher growth of microbial pathogens of soft-rot at 30°C, and storage temperatures ranging from 30 to 35°C as the most favourable for their development on cabbage. Also, Farrar *et al.* (2000) reported temperatures between 30 and 37°C as optimum for soft-rot development in different vegetable crop species. However, Raju *et al.* (2008) observed enhancement of soft-rotting ability of the bacterial *Erwinia carotovora* subsp. *carotovora* at much lower temperatures between 20 and 30°C, with severity of soft-rotting increasing with higher relative humidity. The highest extent of soft-rotting was recorded when samples were incubated at 35°C and 100% relative humidity. At the other extreme, soft-rot diseased tubers placed in dry atmospheric conditions became dehydrated quickly thus, halting the progression of the disease (Walker, 2004).

In Minna, the heavy postharvest losses of sweet potato have been attributed to poor prevailing storage environmental conditions. Yet, information on the influence of environmental factors on the activities of sweet potato soft-rot microbial agents is very scanty and, perhaps, explains the delay in the development of cost-effective postharvest storage strategies for the crop in the country. This study was, therefore, carried out to determine the influence of temperature and relative humidity on growth activities of fungi associated with sweet potato soft-rotting.

MATERIALS AND METHODS

Collection of Sweet Potato Tubers

Samples of Soft-rot diseased sweet potato tubers were collected from different

locations within Minna Emirate, Nigeria. Collected tubers were individually preserved in labeled sterile polythene bags before transportation to the Laboratory for analyses.

Isolation of Micro-organisms

The isolation of fungal microbes associated with soft-rot in the diseased-tubers was carried out following standard procedures. The tubers were first washed under running tap water and then surface sterilized with 70% alcohol. Thereafter, they were cut into 2x2x2mm cubes, under sterile conditions, and each piece of tuber cube was aseptically placed on potato dextrose agar (PDA) medium. The inoculated plates were incubated at room temperature (27±3°C). The fungal colonies that developed were subcultured onto PDA, to obtain pure cultures.

Identification of Isolated Micro-organisms

The fungal isolates were identified using morphological characteristics according to the methods of Barrow and Felthan (1992) and Garrette (1978).

Effect of Temperature on Growth of Fungal Isolates

Pure cultures of the fungal isolates were subsequently inoculated onto PDA media and incubated at five temperature regimen namely, 27±3°C (i.e., room temperature, 'Control'), 35, 40, 45 and 50°C. Each temperature treatment had three replicates and the whole experiment was allowed to stand for eight days. Growth extent of the isolates was estimated at 24-hr intervals, using the Linear Measurement technique.

Effect of Relative Humidity on Growth of Fungal Isolates

Inoculated plates of the fungal isolates were placed individually on copper wire gauze in the upper chamber of desiccators with the lower chamber containing 250ml saturated solution each of the following salts:

Magnesium chloride, Calcium nitrate, Ammonium sulphate, Potassium nitrate, Potassium sulphate and distilled water (i.e., 'Control'), to give relative humidity of 32.5, 50, 74, 85, 97.5 and 100%, respectively. Also, each R.H. treatment had three replicates and the whole experiment was monitored for fungal growth at 24-hr interval for eight days at room temperature (i.e., $27.00 \pm 3.00^\circ\text{C}$).

Data Analysis

Mean daily growth of the fungal isolates at different temperature and R.H. treatments were calculated. For each parameter (i.e., temperature and R.H.), differences in the mean daily growth values for each fungal isolates, among the different treatments were compared for statistical significance using the Chi-square tested, at $P = 0.05$ level of significance.

RESULTS

The influence of temperature on growth of the fungal isolates is presented in Figure 1. Generally, growth activities of the isolates increased till 40.00°C , after which significant ($P < 0.05$) drop in growth was recorded at higher temperatures tested. However, few exceptions to this general observation were recorded. For example, while the growth activity of *Fusarium oxysporum* dropped from 55.29mm at $27 \pm 3^\circ\text{C}$ to 48.75mm at 35.00°C ; *Scopulariopsis brevicaulus* achieved its highest growth at $27 \pm 3^\circ\text{C}$. Also, significant differences in mean weekly growth of the isolates were limited to the first three temperature regimens tested (range = 27 ± 3 to 40°C), while such differences at 45°C were not significant. At

the highest temperature investigated, the fungal isolates had more-or-less equally very poor growth, with the exception of *Aspergillus niger*.

The growth of all fungal isolate equally responded significantly ($P < 0.05$) to increase in Relative Humidity (R.H.), though with different patterns of such responses. While all their fungi recorded their least mean growth at 32.50% R.H. (range = $6.25 \pm 1.39\text{mm}$ in *Trichoderma viridae* to $28.25 \pm 4.36\text{mm}$ in *Mucor racemosus*), *A. niger*, *R. oryzae* and *M. racemosus* achieved maximum growth at 50 R.H. (i.e., 46 ± 23.07 , 53.38 ± 23.37 and $59.63 \pm 28.09\text{mm}$, respectively). On the other hand, *F. oxysporum* and *S. brevicaulus* grew optimally at 85.00% R.H. ($56.43 \pm 28.83\text{mm}$ and $28.38 \pm 18.02\text{mm}$, respectively). Distinctly, the growth of *T. viridae* were highest at the last three R.H. (i.e., 85.00 - 100.00%), with insignificantly different ($P > 0.05$) growth rate (range = $21.00 \pm 14.50\text{mm}$ at 97.50% R.H. to 21.25mm at 85.00 and 100.00% R.H.). On the whole, three of the fungi namely, *F. oxysporum*, *S. brevicaulus* and *T. viridae* continued to grow very well at extremely high R.H. of 97.50 and 100.00%. At the other extreme, i.e., low R.H. of 32.50%, the growth of *S. brevicaulus* and *T. viridae* were seriously impeded (i.e., 8.50 ± 2.07 and $6.25 \pm 1.39\text{mm}$, respectively), while the other fungi achieved more than 20mm mean growth.




Figure 1: Influence of Temperature (°C) on mean diameter of daily growth (mm) of soft-rot fungal isolates from diseased sweet potato tubers

Table 1: Influence of relative humidity (%) on mean diameter of daily growth (mm) of soft-rot fungal isolates from diseased sweet potato tubers

Test Isolates	Mean Growth of Fungi at Different Relative Humidities (%)					
	32.5	50.00	74.60	85.00	97.50	100.00
<i>Aspergillus niger</i>	25.25±15.69 ^{a*}	46.00±23.07 ^c	36.13±37.31 ^b	27.63±35.31 ^a	28.00±36.08 ^a	29.00±37.50 ^a
<i>Fusarium oxysporum</i>	25.13±14.69 ^a	39.13±22.48 ^b	47.50±32.70 ^c	56.43±28.83 ^d	53.86±27.93 ^c	46.75±32.27 ^c
<i>Rhizopus oryzae</i>	22.50±14.05 ^a	53.38±23.37 ^b	23.13±36.54 ^a	24.00±37.50 ^a	23.63±36.15 ^a	23.63±36.15 ^a
<i>Mucor racemosus</i>	28.25±14.36 ^a	59.63±28.09 ^b	23.50±36.12 ^a	24.50±36.83 ^a	24.25±36.75 ^a	24.25±36.75 ^a
<i>Scopulariopsis brevicaulus</i>	8.50±2.07 ^a	20.75±14.35 ^b	25.88±16.98 ^b	28.38±18.02 ^c	27.50±17.35 ^c	27.34±17.35 ^c
<i>Trichoderma viridae</i>	6.25±1.39 ^a	14.53±8.07 ^b	18.25±11.65 ^b	21.25±13.98 ^c	21.00±14.50 ^c	21.25±15.00 ^c

*Values followed by same superscript alphabet in a row are not significantly different at P = 0.05 level of significance

DISCUSSION

To some extent the mean growth of the fungi increased with increasing temperature but succumbed at extremely high temperatures. For most of the species, the optimum temperature was 40°C. This finding confirms the popular belief that temperature is one of the major environmental factors influencing rot at temperatures ranging from 30 – 37°C. Even temperatures as low as 20 – 30°C were

the progression of sweet potato soft-rot disease during storage (Brown and Nelson, 2003). Notably, the optimum temperature for the growth of most of the soft-rot fungi recorded in this study (i.e., 40°C) was relatively higher than reported elsewhere. Bhat *et al.* (2010a&b) and Farrar *et al.* (2000) observed optimal growth of microbial pathogens of soft-reported as optimally enhancing the soft-rotting ability of microbial pathogens.

Usually, the optimum temperature for the growth of microbial pathogens is often dictated by the prevailing temperature of the local environment. This, perhaps, explains the relatively high temperature for the optimum growth of the fungi. The mean temperature during the dry season in the study area, where postharvest storage of sweet potato is practiced, was about 40°C (Olayemi *et al.*, 2011).

The growth of the fungi also responded significantly to increase in Relative Humidity. However, all species had their significantly least weekly growth at the lowest R.H. (i.e., 32.50%). This is to be expected, as microbial pathogens generally require a minimum level of moisture condition to meet their enzymatic and metabolic activities. The results of this study showed that *F. oxysporum*, *T. viridae* and *S. brevicaulus* grew optimally at very high R.H. (i.e., between 85 and 100%), been consistent with those of Raju *et al.* (2008). However, as noticed with the influence of temperature on growth of the fungal isolates in this study, many of the fungi achieved their optimum growth at relative humidities well outside the range generally regarded as conventional but closely related to that prevailing in the locality. In this study, *A. niger*, *R. oryzae* and *M. racemosus* achieved maximum growth at a low R.H. of 50%. However, the annual mean R.H. in Minna was recently reported to be 47.05±21.66% (Olayemi *et al.*, 2011) thus, further buttressing the earlier submission that the sweet potato soft-rot fungal pathogens in Minna have evolved and adapted to the prevailing temperature and R.H. in the area.

CONCLUSION

The growth of sweet potato soft-rot fungal isolates in Minna Emirate were significantly influenced by varying temperature and relative humidity. However, while some of the fungi species grew optimally at

temperature and R.H. considered conventional for soft-rotting microbial pathogens, quite a number of the fungi species achieved optimum growth at temperatures and relative humidities clearly outside the normal range, but closely related to prevailing environmental climatic conditions in Minna thus, suggesting environmentally-induced adaptation of the fungal species. The findings of this study should, therefore, guide the development of better postharvest sweet potato storage facilities, based on appropriate modification of the micro-climatic conditions in Minna Emirate.

REFERENCES

- Abubaker, A. S. (1990). *The sweet potato crop in Kenya*. In: Allard, G. B.; Rangi, D. K. and Abubaker, A. S. (Eds.). Proceedings of Sweet potato Management in Kenya National Workshop, Mombasa, Kenya.
- Barrow, S. and Felthan, W. (1992). *Laboratory manual for identification of bacteria*. 3rd ed. Cambridge University Press, London, England. Pp 61-108.
- Bhat, K. A., Masood, S. D., Bhat, N. A., Ashraf-Bhat, M., Razvi, S. M., Mir, M. R., Akhtar, S., Wani, N. and Habib, M. (2010b). Current status of postharvest soft rot in vegetables: A review. *Asian Journal of Plant Sciences*, ISSN 1682-3974.
- Bhat, K. A., Masoodi, S. D., Bhat, N. A., Ahmad, M., Zargar, M. Y., Mir, S. A. and Ashraf-Bhat, M. (2010a). Studies on the effect of temperature on the development of soft rot of cabbage (*Brassica oleracea var capitata*) caused by *Erwinia carotovora subsp.* *Journal of Phytopathology*, 2: 64-67.
- Brown, C. and Nelson, B. C. (2003). *Storage of potatoes: Postharvest behavior, store design, storage practice, handling*. Center for

Agricultural Publishing and Documentation, Wageningen.

FAO. (1992). *The world sweet potato economy*. Food and Agricultural Organisation. Rome, Italy.

Farrar, J. J., Nunez, J. J. and Davis, R. M. (2000). Influence of soil saturation and temperature on *Erwinia chrysanthemi* causing soft rot of carrot. *Plant Diseases*, 84: 665-668.

Garrette, S. D. (1978). *Plant pathology: Principles and practice*. Open University Press, Milton Keynes, pp 237.

George, J. B. (1988). Postharvest handling, transporting and losses during marketing of sweet potato. *Acta Horticulturae*, 218:369-373.

Kihurani, A. W., Narla, R. D., Shibairo, S., Imungi, J. and Carey, E. (2008). Effect of soil pH on postharvest pathological deterioration of sweet potato storage roots. *African Journal of Horticultural Science*, 1:1-8.

Mutuura, J. N., Ewell, P. T., Abubakar, A., Munga, T., Ajanga, S., Irungu, J., Omari, F. and Maobe, S. (1992). Sweet potato in the food systems of Kenya: Results of a socio-

economic survey. *In*: Kabira, J. N. and Ewell, P. T. (Eds.). *Current Research for the Improvement of Potato and Sweet potato in Kenya*. Proceedings of KARI/CIP Technical Workshop on Collaborative Research, Nairobi, Kenya.

Olayemi, I. K., Ande, A. T., Ayanwale, A. V., Mohammed, A. Z., Bello, I. M., Idris, B., Isah, B., Chukwuemeka, V. and Ukubuiwe, A. C. (2011). Seasonal trends in epidemiological and entomological profiles of malaria transmission in north central Nigeria. *Pakistan Journal of Biological Sciences*, 14(4): 293-299.

Raju, M. R. B., Pal, V. and Jalali, I. (2008). (2008). Inoculation method of *Pectobacterium carotovorum sub sp. Carotovorum* and factors influencing development of bacterial soft rot in Radish. *Journal of Mycology and Plant Pathology*, 38: 311-315.

Walker, J. C. (1998). *Bacterial soft rots of carrot*. Discovery Publishing House, New Delhi, pp. 78.

Walker, J. C. (2004). Bacterial soft rot. *In: Manual of vegetable plant diseases*, Chupp, C. (Ed.). Shristi Book Distributors, New Delhi.